

SUSY effects on the exclusive semi-leptonic decays $B \rightarrow \pi\tau^+\tau^-$ and $B \rightarrow \rho\tau^+\tau^-$

S.Rai Choudhury* and Naveen Gaur†

Department of Physics & Astrophysics,

University of Delhi,

Delhi - 110 007, India

(Dated: February 1, 2008)

Semi-leptonic decays of B-meson are known to be very sensitive to the any new physics effects. Amongst various possibilities the transition at the quark level $b \rightarrow d\ell^+\ell^-$ is more suited than $b \rightarrow s\ell^+\ell^-$ for purpose of study of CP violation. Here in this work we will discuss the effects of Supersymmetry on the various experimentally measurable quantities like decay rate, Forward backward asymmetry, various polarization asymmetries and CP violation asymmetries in exclusive channels $B \rightarrow \pi\ell^+\ell^-$ and $B \rightarrow \rho\ell^+\ell^-$. We will be going to focus mainly on the Neutral Higgs boson effects on these measurements, with a view of eliciting information about possible CP-violating as well as non-hermitian terms in the effective Hamiltonian.

PACS numbers: 13.20.He, 12.60.Jv, 13.88.+e

I. INTRODUCTION

The rare B meson decays induced by Flavor changing neutral current (FCNC) $b \rightarrow s(d)$ transition offer a deeper probe for weak interaction sector of the Standard model (SM) as they go through second order in weak interaction. These decays can give information regarding the fundamental parameters like Cabbibo-Kobayashi-Maskawa (CKM) factors, leptonic decay constants etc. These decays can also be very useful in testing out the various new physics scenarios like 2HDM models [2, 3], MSSM [4, 5, 6, 10, 11]. Among the weak decays of B-meson the leptonic and semi-leptonic decays are very useful because of their relative cleanness. Among these too the exclusive decays lately have received special attention [8, 9]. For the semi-leptonic and leptonic decays like $B \rightarrow X_{s,d}\ell^+\ell^-$, $B \rightarrow K(K^*)\ell^+\ell^-$, $B \rightarrow \pi\ell^+\ell^-$, $B \rightarrow \rho\ell^+\ell^-$ etc. the basic quark level process is $b \rightarrow s(d)\ell^+\ell^-$ [1]. The basic quark level process $b \rightarrow s(d)\ell^+\ell^-$ occurs through intermediate t, c or u quark. These processes can be described in terms of effective Hamiltonian which contains the information about short and long distance effects. For the quark level process $b \rightarrow s\ell^+\ell^-$ the various contributions due to intermediate t, c and u quarks enters in matrix elements with factors $V_{tb}V_{ts}^*$, $V_{cb}V_{cs}^*$ and $V_{ub}V_{us}^*$. Off these three factors the third one is extremely small as compared to the first two. The unitarity relationship of CKM matrix becomes (approximately) $V_{tb}V_{ts}^* + V_{cb}V_{cs}^* \approx 0$, so that the second factor can be written effectively as the negative of the first one. The Hamiltonian (effective) for $b \rightarrow s\ell^+\ell^-$ transition thus involves essentially only one independent CKM factor $V_{tb}V_{ts}^*$ and hence the process $b \rightarrow s\ell^+\ell^-$ is not sensitive to CKM phases within SM. [12].

For the transition $b \rightarrow d\ell^+\ell^-$ the CKM factors involved $V_{tb}V_{td}^*$, $V_{cb}V_{cd}^*$ and $V_{ub}V_{ud}^*$ are comparable in magnitude and so the crossection for the processes having the quark level process $b \rightarrow d\ell^+\ell^-$ can have significant interference terms between them, and this could open up the possibility of observing the complex CKM factors from the interference

*Electronic address: src@ducos.ernet.in

†Electronic address: naveen@physics.du.ac.in; URL: <http://physics.du.ac.in/~naveen>

terms. In the semi-leptonic decays (having lepton pair in final state) one can discuss several other kinematical variables associated with final state leptons, like lepton pair Forward Backward (FB) asymmetry and various polarization asymmetries. SUSY effects on the FB asymmetries in various exclusive decay modes of B-meson, like $B \rightarrow (\pi, K)\ell^+\ell^-$ [8], $B \rightarrow (\pi, \rho)\ell^+\ell^-$ [8, 9], $B \rightarrow \ell^+\ell^-\gamma$ [7] have been extensively studied. Sometime back as pointed out by one of us [13] for inclusive decay mode $B \rightarrow X_d\ell^+\ell^-$ and by Krüger and Sehgal [14] for exclusive mode $B \rightarrow (\pi, \rho)\ell^+\ell^-$ that along with CP violation from partial width asymmetry one can look for CP violation in FB asymmetry also, we will be going to explore this possibility also.

Along with FB asymmetry associated with final state leptons one can also discuss the three polarization asymmetries (longitudinal, normal and transverse) associated with final state leptons in various semi-leptonic decays. The importance of polarization asymmetries associated with final state leptons in various inclusive and exclusive semi-leptonic decay modes have been extensively discussed in many works [6, 7, 15, 16] . In this communication, we study the three possible polarization asymmetries with the idea of exploring CP asymmetries as well as the *non hermiticity* of the effective Hamiltonian¹ through their measurement.

The paper is organized as follows : In section II we will be going to present the general formalism where we will be going to write the general effective Hamiltonian and will be going to present our definitions of the FB asymmetry and polarization asymmetries. Section III will be devoted to the decay mode $B \rightarrow \pi\ell^+\ell^-$ where we will discuss all the kinematical variables associated with this decay mode. In Section IV we will discuss the decay mode $B \rightarrow \rho\ell^+\ell^-$. Finally we will conclude with results and discussion in section V.

II. GENERAL FORMALISM

The QCD corrected effective Hamiltonian for the decay $b \rightarrow d\ell^+\ell^-$ in general SUSY model can be written as [5] :

$$\mathcal{H}_{eff} = \frac{4G_F\alpha}{\sqrt{2}\pi} V_{tb}V_{td}^* \left[\sum_{i=1}^{10} C_i O_i + \sum_{i=1}^{10} C_{Q_i} Q_i - \lambda_u \{ C_1 [O_1^u - O_1] + C_2 [O_2^u - O_2] \} \right], \quad (2.1)$$

where we have used the unitarity of the CKM matrix $V_{tb}V_{td}^* + V_{ub}V_{ud}^* \approx -V_{cb}V_{cd}^*$, and $\lambda_u = V_{ub}V_{ud}^*/V_{tb}V_{td}^*$. Here O_1 and O_2 are the current current operators, $O_3, \dots, O_6$ are called QCD penguin operators and O_9 and O_{10} are semileptonic electroweak penguin operators [1]. The new operators Q_i ($i = 1, \dots, 10$) arises due to NHB (Neutral Higgs Boson) exchange diagrams [3, 5]. In addition to the short distance corrections included in the Wilson coefficients, there are some long distance effects also, associated with real $c\bar{c}$ resonances in the intermediate states. This is taken into account by using the prescription given in [20] namely by using the Breit-Wigner form of the resonances that adds to C_9^{eff} :

$$C_9^{res} = \frac{-3\pi}{\alpha^2} \kappa_v \sum_{V=J/\psi, \psi', \dots} \frac{M_V Br(V \rightarrow \ell^+\ell^-) \Gamma_{total}^V}{(s - M_V^2) + i\Gamma_{total}^V M_V}, \quad (2.2)$$

as we are taking final leptons to be τ so only five resonances of the $c\bar{c}$ will contribute. The phenomenological factor κ_v is taken to be 2.3 for numerical calculations [16] In this work we will be going to use the Wolfenstein parameterisation [22] of CKM matrix with four real parameters λ, A, ρ and η where η is the measure of CP violation. In terms of these

¹ which arises out of the quark loops

parameters we can write λ_u as :

$$\lambda_u = \frac{\rho(1-\rho) - \eta^2}{(1-\rho)^2 + \eta^2} - i \frac{\eta}{(1-\rho)^2 + \eta^2} + O(\lambda^2), \quad (2.3)$$

From the relevant part of the above effective Hamiltonian eqn.(2.1) we can write the QCD corrected matrix element as :

$$\begin{aligned} \mathcal{M} = \frac{G_F \alpha}{\sqrt{2}\pi} V_{tb} V_{td}^* \left\{ -2 C_7^{eff} \frac{m_b}{q^2} (\bar{d} i \sigma_{\mu\nu} q^\nu P_R b) (\bar{\ell} \gamma^\mu \ell) + C_9^{eff} (\bar{d} \gamma_\mu P_L b) (\bar{\ell} \gamma^\mu \ell) + C_{10} (\bar{d} \gamma_\mu P_L b) (\bar{\ell} \gamma^\mu \ell) \right. \\ \left. + C_{Q_1} (\bar{d} P_R b) (\bar{\ell} \ell) + C_{Q_2} (\bar{d} P_R b) (\bar{\ell} \gamma_5 \ell) \right\}, \end{aligned} \quad (2.4)$$

where q is the momentum transfer and $P_{L,R} = (1 \mp \gamma_5)/2$, we have neglected mass of d quark. The wilson coefficients C_7^{eff} and C_{10} are given in many works [5, 11, 19] and the other Wilsons C_{Q_1} and C_{Q_2} are given in [4, 5]. The definition of C_9^{eff} is given in [9, 13, 17].

The decay rate (for any general three body decay process $B \rightarrow P \ell^+ \ell^-$) can be evaluated by doing the phase space integration. On doing the phase space integration we can get

$$\frac{d\Gamma(B \rightarrow P \ell^+ \ell^-)}{d\hat{s} dx} = \frac{m_B}{2^9 \pi^3} \lambda^{1/2}(1, \hat{s}, \hat{m}_p^2) \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} |\mathcal{M}|^2, \quad (2.5)$$

where $\hat{s} = s/m_B^2$, $\hat{m}_\ell = m_\ell/m_B$, $\hat{m}_p = m_p/m_B$ are the dimensionless quantities. $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2ab - 2ac - 2bc$. s is the CM energy of $\ell^+ \ell^-$ system, m_P is the mass of particle labeled as P and $z = \cos\theta$ where θ is the angle between ℓ^- and B three momenta in CM frame of $\ell^+ \ell^-$. $|\mathcal{M}|^2$ is the matrix element square of the process under consideration.

From above expression one can get decayrate, FB asymmetry [20]. Decay rate is simply the integration of eqn(2.5) over the angle z . The definition of the FB asymmetry is [20]:

$$A_{FB} = \frac{\int_0^1 dz \frac{d\Gamma}{d\hat{s} dz} - \int_{-1}^0 dz \frac{d\Gamma}{d\hat{s} dz}}{\int_0^1 dz \frac{d\Gamma}{d\hat{s} dz} + \int_{-1}^0 dz \frac{d\Gamma}{d\hat{s} dz}}, \quad (2.6)$$

For defining the polarization asymmetries we define the orthogonal unit vectors, S in the rest frame of ℓ^- for the polarization of lepton ℓ^- [6, 16] to the longitudinal direction (L), the normal direction (N) and the transverse direction (T).

$$\begin{aligned} S_L^\mu &\equiv (0, \mathbf{e}_L) = (0, \frac{\mathbf{p}_-}{|\mathbf{p}_-|}) \\ S_N^\mu &\equiv (0, \mathbf{e}_N) = (0, \frac{\mathbf{q} \times \mathbf{p}_-}{|\mathbf{q} \times \mathbf{p}_-|}) \\ S_T^\mu &\equiv (0, \mathbf{e}_T) = (0, \mathbf{e}_N \times \mathbf{e}_L) \end{aligned} \quad (2.7)$$

where $\mathbf{p}_-$ and $\mathbf{q}$ are the three momenta of ℓ^- and photon in the center-of-mass (CM) frame of $\ell^- \ell^+$ system. Now boosting all the three vectors given in eqn.(2.7), only longitudinal vector will get boosted and rest two (normal and transverse) will remain the same. Longitudinal vector after boost becomes

$$S_L^\mu = (\frac{|\mathbf{p}_-|}{m_\ell}, \frac{E_- \mathbf{p}_-}{m_\ell |\mathbf{p}_-|}) \quad (2.8)$$

Now we can calculate the polarization asymmetries by using the spin projectors for ℓ^- as $\frac{1}{2}(1 + \gamma_5 \not{S})$. The lepton polarization asymmetries are defined as :

$$P_x(\hat{s}) \equiv \frac{\frac{d\Gamma(S_x)}{d\hat{s}} - \frac{d\Gamma(-S_x)}{d\hat{s}}}{\frac{d\Gamma(S_x)}{d\hat{s}} + \frac{d\Gamma(-S_x)}{d\hat{s}}} \quad (2.9)$$

with $x = L, T, N$ respectively for longitudinal, transverse and normal polarization asymmetry.

Now ready with the terminology and definitions we will move to the calculations of the various measurable quantities that we mentioned before.

III. $B \rightarrow \pi \ell^+ \ell^-$ DECAY MODE

In this section we calculate the Branching ratio, FB asymmetry and the polarization asymmetries associated with the inclusive decay mode $B \rightarrow \pi \ell^+ \ell^-$. Using the definition of the form factors (eqn(B1)-(B3)) we can write down the matrix element for $\bar{B} \rightarrow \pi^0$ transition as ³ :

$$\mathcal{M}^{\bar{B} \rightarrow \pi} = \frac{G_F \alpha}{\sqrt{2} \pi} V_{tb} V_{td}^* \{ A(p_B)_\mu (\bar{\ell} \gamma_\mu \ell) + B(p_B)_\mu (\bar{\ell} \gamma_\mu \gamma_5 \ell) + C(\bar{\ell} \gamma_5 \ell) + D(\bar{\ell} \ell) \} \quad (3.1)$$

with

$$A = C_9^{eff} F_1(q^2) - 2C_7^{eff} \tilde{F}_T(q^2) \quad (3.2)$$

$$B = C_{10} F_1(q^2) \quad (3.3)$$

$$C = m_\ell C_{10} \left\{ -F_1(q^2) + \frac{(m_B^2 - m_\pi^2)}{q^2} (F_0(q^2) - F_1(q^2)) \right\} + \frac{(m_B^2 - m_\pi^2)}{2m_b} F_0(q^2) C_{Q_2} \quad (3.4)$$

$$D = \frac{(m_B^2 - m_\pi^2)}{2m_b} F_0(q^2) C_{Q_1} \quad (3.5)$$

From the above expression of the matrix element (eqn.(3.1)) we can get the analytical expression of the decay rate as

$$\frac{d\Gamma(\bar{B} \rightarrow \pi \ell^+ \ell^-)}{d\hat{s}} = \frac{G_F^2 m_B^5 \alpha^2}{3 \times 2^9 \pi^5} |V_{tb} V_{td}^*|^2 \lambda^{1/2}(1, \hat{s}, \hat{m}_\pi^2) \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \Sigma_\pi \quad (3.6)$$

with

$$\begin{aligned} \Sigma_\pi = & \lambda(1, \hat{s}, \hat{m}_\pi^2) \left(1 + \frac{2\hat{m}_\ell^2}{\hat{s}} \right) |A|^2 + \left[\lambda(1, \hat{s}, \hat{m}_\pi^2) \left(1 + \frac{2\hat{m}_\ell^2}{\hat{s}} \right) + 24\hat{m}_\ell^2 \right] |B|^2 + 6 \frac{\hat{s}}{m_B^2} |C|^2 \\ & + 6 \frac{\hat{s}}{m_B^2} \left(1 - \frac{4\hat{m}_\ell^2}{\hat{s}} \right) |D|^2 + 12 \frac{\hat{m}_\ell}{m_B} (1 + \hat{s} - \hat{m}_\pi^2) \text{Re}(C^* B) \end{aligned} \quad (3.7)$$

The expression of FB asymmetry is :

$$A_{FB}(\bar{B} \rightarrow \pi \ell^+ \ell^-) = -6 \hat{m}_\ell \lambda^{1/2}(1, \hat{s}, \hat{m}_\pi^2) \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \frac{\text{Re}(AD^*)}{(m_B \Sigma_\pi)} \quad (3.8)$$

As we can see from above expression that FB asymmetry is proportional to the new interactions , i.e. ,NHB contributions. This is a point which has also been noted in some earlier works [8].

We divide this section in two subsections in the first one we will discuss the CP violation in $B \rightarrow \pi \ell^+ \ell^-$ and in next subsection we will discuss the polarization asymmetries associated with final state leptons.

² $\bar{B}$ actually is B^+

³ in writing this we have used $\bar{\ell} \not{A} \ell = 0$, $\bar{\ell} \not{A} \gamma_5 \ell = 2m_\ell \bar{\ell} \gamma_5 \ell$




FIG. 1: Branching ratio for $\bar{B} \rightarrow \pi \tau^+ \tau^-$. Other parameters are : mSUGRA $m = 200$, $M = 450$, $A = 0$, $\tan \beta = 35$, $\text{sgn}(\mu)$ is taken to be +ve for rSUGRA $m_A = 306$. All masses are in GeV

A. CP violation

Firstly we define CP-violating partial width asymmetry between B and $\bar{B}$ decay. This is defined as :

$$A_{CP}(\hat{s}) = \frac{\frac{d\Gamma}{d\hat{s}} - \frac{d\bar{\Gamma}}{d\hat{s}}}{\frac{d\Gamma}{d\hat{s}} + \frac{d\bar{\Gamma}}{d\hat{s}}} \quad (3.9)$$




FIG. 2: FB asymmetry for $\bar{B} \rightarrow \pi \tau^+ \tau^-$. Other parameters are : mSUGRA $m = 200$, $M = 450$, $A = 0$, $\tan \beta = 35$, $\text{sgn}(\mu)$ is taken to be +ve for rSUGRA $m_A = 306$. All masses are in GeV

where

$$\frac{d\Gamma}{d\hat{s}} = \frac{d\Gamma(\bar{B} \rightarrow \pi \ell^+ \ell^-)}{d\hat{s}} \quad , \quad \frac{d\bar{\Gamma}}{d\hat{s}} = \frac{d\Gamma(B \rightarrow \bar{\pi} \ell^+ \ell^-)}{d\hat{s}} \quad (3.10)$$

In going from Γ to $\bar{\Gamma}$ the only change we have to make is in the expression of C_9^{eff} . Defining C_9^{eff} as :

$$C_9^{eff} = \xi_1 + \lambda_u \xi_2 \quad (3.11)$$

where ξ_1 , ξ_2 and λ_u all are complex. In going from $\bar{B} \rightarrow \pi \ell^+ \ell^-$ to $B \rightarrow \bar{\pi} \ell^+ \ell^-$, C_9^{eff} becomes :

$$C_9^{eff} = \xi_1 + \lambda_u^* \xi_2 \quad (3.12)$$

with this change one can get the expression for $d\bar{\Gamma}/d\hat{s}$ as :

$$\frac{d\Gamma(B \rightarrow \bar{\pi} \ell^+ \ell^-)}{d\hat{s}} = \frac{G_F^2 m_B^5 \alpha^2}{3 \times 2^9 \pi^5} |V_{tb} V_{td}^*|^2 \lambda^{1/2}(1, \hat{s}, \hat{m}_\pi^2) \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \{ \Sigma_\pi + 4 \operatorname{Im} \lambda_u \Delta_\pi \} \quad (3.13)$$

with

$$\Delta_\pi = \left\{ \operatorname{Im}(\xi_1^* \xi_2) |F_1(s)|^2 - 2 C_7^{eff} \operatorname{Im} \xi_2 F_T(s) F_1(s) \frac{m_b}{(m_B + m_\pi)} \right\} \lambda(1, \hat{s}, \hat{m}_\pi^2) \left(1 + \frac{2\hat{m}_\ell^2}{\hat{s}} \right) \quad (3.14)$$

Using eqn.(3.6),(3.13) and eqn.(3.14) we can get the CP violating partial width asymmetry as :

$$A_{CP}(\hat{s}) = \frac{-2 \operatorname{Im} \lambda_u \Delta_\pi}{\Sigma_\pi + 2 \operatorname{Im} \lambda_u \Delta_\pi} \quad (3.15)$$




FIG. 3: CP violating asymmetry (A_{CP}) in $\bar{B} \rightarrow \pi \tau^+ \tau^-$ and $B \rightarrow \bar{\pi} \tau^+ \tau^-$. The Wolfenstein parameter we have choosen are : $\rho = -0.07$, $\eta = 0.34$. Other parameters are : mSUGRA $m = 200$, $M = 450$, $A = 0$, $\tan \beta = 35$, $\operatorname{sgn}(\mu)$ is taken to be +ve for rSUGRA $m_A = 306$. All masses are in GeV

As argued in earlier work [13] that by measuring the FB asymmetries of B and $\bar{B}$ one can observe the CP violating phase of CKM matrix. We will here give as to how much the predictions of SM would change in SUSY.

In the discussion of CP violation by measuring FB asymmetry its important to fix the sign of convention. The sign of FB asymmetry for B and $\bar{B}$ decays are different. In fact under strict CP conservation :

$$A_{FB}(\bar{B}) = - A_{FB}(B) \quad (3.16)$$

So under CP conservation the FB asymmetry of B and $\bar{B}$ are exactly opposite ⁴. So any change in eqn.(3.15) will be measure of CP violation. We define CP violating parameter of FB asymmetry as :

$$\delta_{FB} = A_{FB}(\bar{B} \rightarrow \pi \ell^+ \ell^-) + A_{FB}(B \rightarrow \bar{\pi} \ell^+ \ell^-) \quad (3.17)$$

We can get the expression of δ_{FB} from the expression of FB asymmetry eqn.(3.8) as :

$$\begin{aligned} \delta_{FB} = & \frac{12\hat{m}_\ell \lambda^{1/2}(1, \hat{s}, \hat{m}_\pi^2) \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}}}{m_B \Sigma_\pi (\Sigma_\pi + 4Im\lambda_u \Delta_\pi)} Im\lambda_u \left[\Sigma_\pi F_1(s) Im\xi_2 \right. \\ & \left. + \Delta_\pi (2C_7^{eff} \tilde{F}_T(s) + F_1(s) + Im\xi_2 Im\lambda_u - F_1(s) Re\xi_1 - F_1(s) Re\xi_2 Re\lambda_u) \right] \end{aligned} \quad (3.18)$$




FIG. 4: CP violating asymmetry (δ_{CP}) in $\bar{B} \rightarrow \pi \tau^+ \tau^-$ and $B \rightarrow \bar{\pi} \tau^+ \tau^-$. The Wolfenstein parameter we have choosen are : $\rho = -0.07$, $\eta = 0.34$. Other parameters are : mSUGRA $m = 200$, $M = 450$, $A = 0$, $\tan \beta = 35$, $sgn(\mu)$ is taken to be +ve for rSUGRA $m_A = 306$. All masses are in GeV

⁴ we can understand this -ve sign because under CP conjugation not only $b \leftrightarrow \bar{b}$ but there is a transformation in leptons also and $\ell^+ \leftrightarrow \ell^-$. Since the two leptons are emitted back to back in cm frame of dileptons the FB asymmetry defined in terms of negatively charged lepton, ℓ^- (for both B and $\bar{B}$), this changes sign under CP conjugation

B. Polarization asymmetries

We can also get the expression of the polarization asymmetries in $\bar{B} \rightarrow \pi \ell^+ \ell^-$ transtion using the formalism given in section II. The expression of the various polarization asymmetries of ℓ^- are :

$$P_L = 0 \quad (3.19)$$

$$P_T = 0 \quad (3.20)$$

$$P_N = \frac{\pi D \lambda^{1/2}(1, \hat{s}, \hat{m}_\pi^2) \sqrt{\hat{s} - 4\hat{m}_\ell^2}}{\Sigma_\pi} \left[Im\xi_1 + Im\lambda_u Re\xi_2 + Im\xi_2 Re\lambda_u \right] \quad (3.21)$$




FIG. 5: Normal polarization asymmetry (P_N) in $\bar{B} \rightarrow \pi \tau^+ \tau^-$. The Wolfenstein parameter we have choosen are : $\rho = -0.07$, $\eta = 0.34$. Other parameters are : mSUGRA $m = 200$, $M = 450$, $A = 0$, $\tan\beta = 35$, $sgn(\mu)$ is taken to be +ve for rSUGRA $m_A = 306$. All masses are in GeV

IV. $B \rightarrow \rho \ell^+ \ell^-$ DECAY MODE

In this section we will calculate the possible measurables associated with the inclusive decay mode $B \rightarrow \rho \ell^+ \ell^-$. Using the defination of the form factors for $B \rightarrow \rho$ transition given by eqns(C1),(C2),(C3) we can write down the matrix elment as :

$$\begin{aligned} \mathcal{M}^{\bar{B} \rightarrow \rho} = & \left[i\epsilon_{\mu\nu\alpha\beta} \epsilon^{\nu*} p_B^\beta q^\beta A + \epsilon_\mu^* B + (\epsilon^* \cdot q)(p_B)_\mu C \right] (\bar{\ell} \gamma^\mu \ell) \\ & + \left[i\epsilon_{\mu\nu\alpha\beta} \epsilon^{\nu*} p_B^\alpha q^\beta D + \epsilon_\mu^* E + (\epsilon^* \cdot q)(p_B)_\mu F \right] (\bar{\ell} \gamma^\mu \ell) + G (\epsilon^* \cdot q)(\bar{\ell} \ell) + H (\epsilon^* \cdot q)(\bar{\ell} \gamma_5 \ell) \end{aligned} \quad (4.1)$$

where

$$A = 4 \frac{C_7^{eff}}{s} m_b T_1(s) + C_9^{eff} \frac{V(s)}{(m_B + m_\rho)} \quad (4.2)$$

$$B = -2 \frac{C_7^{eff}}{s} m_b (m_B^2 - m_\rho^2) T_2(s) - \frac{1}{2} (m_B + m_\rho) A_1(s) C_9^{eff} \quad (4.3)$$

$$C = 4 \frac{C_7^{eff}}{s} m_b \left\{ T_2(s) + \frac{s}{(m_B^2 - m_\rho^2)} T_3(\hat{s}) \right\} + C_9^{eff} \frac{A_2(s)}{m_B + m_\rho} \quad (4.4)$$

$$D = C_{10} \frac{V(s)}{m_B + m_\rho} \quad (4.5)$$

$$E = -\frac{1}{2} (m_B + m_\rho) A_1(s) \quad (4.6)$$

$$F = C_{10} \frac{A_2(s)}{m_B + m_\rho} \quad (4.7)$$

$$G = -C_{Q_1} \frac{m_\rho A_0(s)}{m_b} \quad (4.8)$$

$$H = -C_{Q_2} \frac{m_\rho A_0(s)}{m_b} - C_{10} \frac{m_\ell A_2(s)}{m_B + m_\rho} + \frac{2m_\rho m_\ell}{s} (A_3(s) - A_0(s)) C_{10} \quad (4.9)$$

from the above expression of the matrix element we can get the expression of the partial decay rate :

$$\frac{d\Gamma(\bar{B} \rightarrow \rho \ell^+ \ell^-)}{d\hat{s}} = \frac{G_F^2 m_B^5 \alpha^2}{3 \times 2^{10} \pi^5} |V_{tb} V_{td}^*|^2 \lambda^{1/2}(1, \hat{s}, \hat{m}_\rho^2) \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \Sigma_\rho \quad (4.10)$$

with

$$\begin{aligned} \Sigma_\rho = & (1 + \frac{2\hat{m}_\ell^2}{\hat{s}}) \lambda(1, \hat{s}, \hat{m}_\rho^2) \left[4 m_B^2 \hat{s} |A|^2 + \frac{2}{m_B^2 \hat{m}_\rho^2} (1 + 12 \frac{\hat{m}_\rho^2 \hat{s}}{\lambda(1, \hat{s}, \hat{m}_\rho^2)}) |B|^2 \right. \\ & + \frac{m_B^2}{2\hat{m}_\rho^2} \lambda(1, \hat{s}, \hat{m}_\rho^2) |C|^2 + \frac{2}{\hat{m}_\rho^2} (1 - \hat{m}_\rho^2 + \hat{s}) \text{Re}(B^* C) \left. \right] + 4 m_B^2 \lambda(1, \hat{s}, \hat{m}_\rho^2) (\hat{s} - 4\hat{m}_\ell^2) |D|^2 \\ & + \frac{2}{m_B^2} \left[\frac{2(2\hat{m}_\ell^2 + \hat{s}) - 2(2\hat{m}_\ell^2 + \hat{s})(\hat{m}_\rho^2 + \hat{s}) + 2\hat{m}_\ell^2(\hat{m}_\rho^4 - 26\hat{m}_\rho^2 + \hat{s}^2) + \hat{s}(\hat{m}_\rho^4 + 10\hat{m}_\rho^2 \hat{s} + \hat{s}^2)}{\hat{m}_\rho^2 \hat{s}} \right] |E|^2 \\ & + \frac{m_B^2}{2\hat{m}_\rho^2 \hat{s}} \lambda(1, \hat{s}, \hat{m}_\rho^2) \left[(2\hat{m}_\ell^2 + \hat{s})(\lambda(1, \hat{s}, \hat{m}_\rho^2) + 2\hat{s} + 2\hat{m}_\rho^2) - 2 \{ 2\hat{m}_\ell^2(\hat{m}_\rho^2 - 5\hat{s}) + \hat{s}(\hat{m}_\rho^2 + \hat{s}) \} \right] |F|^2 \\ & - 3 \frac{4\hat{m}_\ell^2 - \hat{s}}{\hat{m}_\rho^2} \lambda(1, \hat{s}, \hat{m}_\rho^2) |G|^2 + 3 \frac{\hat{s}}{\hat{m}_\rho^2} \lambda(1, \hat{s}, \hat{m}_\rho^2) |H|^2 + \frac{2\lambda(1, \hat{s}, \hat{m}_\rho^2)}{\hat{m}_\rho^2 \hat{s}} \left[-2\hat{m}_\ell^2(\hat{m}_\rho^2 - 5\hat{s}) + (2\hat{m}_\ell^2 + \hat{s}) \right. \\ & \left. - \hat{s}(\hat{m}_\rho^2 + \hat{s}) \right] \text{Re}(E^* F) + \frac{12\hat{m}_\ell}{m_B \hat{m}_\rho^2} \lambda(1, \hat{s}, \hat{m}_\rho^2) \text{Re}(H^* E) + \frac{2m_B \hat{m}_\ell}{\hat{m}_\rho^2} \lambda(1, \hat{s}, \hat{m}_\rho^2) (1 - \hat{m}_\rho^2 + \hat{s}) \text{Re}(H^* F) \quad (4.11) \end{aligned}$$

For $B \rightarrow \rho$ transition we can find FB asymmetry to be :

$$\begin{aligned} A_{FB}(\bar{B} \rightarrow \rho \ell^+ \ell^-) = & \left\{ -12\lambda^{1/2}(1, \hat{s}, \hat{m}_\rho^2) \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \hat{s} \left[\text{Re}(A^* D) + \text{Re}(A^* E) \right] - 3 \frac{\hat{m}_\ell \lambda^{1/2}(1, \hat{s}, \hat{m}_\rho^2)}{\hat{m}_\rho^2} \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \right. \\ & \left. \left[2\text{Re}(G^* B) \frac{(1 - \hat{m}_\rho^2 - \hat{s})}{m_B} + \text{Re}(G^* C) m_B \lambda(1, \hat{s}, \hat{m}_\rho^2) \right] \right\} / \Sigma_\rho \quad (4.12) \end{aligned}$$

We will discuss the CP violation in $B \rightarrow \rho$ transition and discuss the polarization asymmetries associated with final state lepton in next subsections.




FIG. 6: Branching ratio for $\bar{B} \rightarrow \rho \tau^+ \tau^-$. Other parameters are : mSUGRA $m = 200$, $M = 450$, $A = 0$, $\tan \beta = 35$, $\text{sgn}(\mu)$ is taken to be +ve for rSUGRA $m_A = 306$. All masses are in GeV

A. CP violation

To find out the CP violating partial width asymmetry we require the expression of the partial width of $B \rightarrow \bar{\rho} \ell^+ \ell^-$. The expression of partial decay rate for $B \rightarrow \bar{\rho} \ell^+ \ell^-$ is :

$$\frac{d\Gamma(B \rightarrow \bar{\rho} \ell^+ \ell^-)}{d\hat{s}} = \frac{G_F^2 m_B^5 \alpha^2}{3 \times 2^{10} \pi^5} |V_{tb} V_{td}^*|^2 \lambda^{1/2}(1, \hat{s}, \hat{m}_\rho^2) \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} (\Sigma_\rho + 4 \text{Im} \lambda_u \Delta_\rho) \quad (4.13)$$




FIG. 7: FB asymmetry for $\bar{B} \rightarrow \rho \tau^+ \tau^-$. Other parameters are : mSUGRA $m = 200$, $M = 450$, $A = 0$, $\tan \beta = 35$, $\text{sgn}(\mu)$ is taken to be +ve for rSUGRA $m_A = 306$. All masses are in GeV




FIG. 8: CP violating asymmetry (A_{CP}) in $\bar{B} \rightarrow \rho\tau^+\tau^-$ and $B \rightarrow \bar{\rho}\tau^+\tau^-$. The Wolfenstein parameter we have chosen are : $\rho = -0.07$, $\eta = 0.34$. Other parameters are : mSUGRA $m = 200$, $M = 450$, $A = 0$, $\tan\beta = 35$, $\text{sgn}(\mu)$ is taken to be +ve for rSUGRA $m_A = 306$. All masses are in GeV

with

$$\begin{aligned} \Delta_\rho = & \left[\text{Im}(\xi_1^* \xi_2) \left\{ 4\hat{s} \frac{|V(s)|^2}{1 + \hat{m}_\rho^2} + (1 + \hat{m}_\rho^2) \left(\frac{6\hat{s}}{\lambda(1, \hat{s}, \hat{m}_\rho^2)} + \frac{1}{2\hat{m}_\rho^2} \right) |A_1(s)|^2 + \frac{\lambda(1, \hat{s}, \hat{m}_\rho^2)}{2\hat{m}_\rho^2(1 + m_\rho)^2} |A_2(s)|^2 \right. \right. \\ & - \left. \frac{1 - \hat{m}_\rho^2 - \hat{s}}{\hat{m}_\rho^2} A_1(s) A_2(s) \right\} + 2 \frac{C_7^{eff} \hat{m}_b}{\hat{s}} \text{Im}(\xi_2) \left\{ 8 \frac{T_1(s) V(s) \hat{s}}{1 + \hat{m}_\rho} \right. \\ & + 2A_1(s) T_2(s) (1 + \hat{m}_\rho)^2 (1 - \hat{m}_\rho) \left(6 \frac{\hat{s}}{\lambda(1, \hat{s}, \hat{m}_\rho^2)} + \frac{1}{2\hat{m}_\rho^2} \right) + A_2(s) \left(T_2(s) + \frac{\hat{s}}{1 - \hat{m}_\rho^2} T_3(s) \right) \frac{\lambda(1, \hat{s}, \hat{m}_\rho^2)}{\hat{m}_\rho^2(1 + \hat{m}_\rho)} \\ & - (1 + \hat{m}_\rho) A_1(s) \left(T_2(s) + \frac{\hat{s}}{1 - \hat{m}_\rho^2} T_3(s) \right) \frac{1 - \hat{m}_\rho^2 - \hat{s}}{\hat{m}_\rho^2} \\ & \left. \left. + A_2(s) T_2(s) (1 - \hat{m}_\rho) \frac{1 - \hat{m}_\rho^2 - \hat{s}}{\hat{m}_\rho^2} \right\} \right] \left(1 + \frac{2\hat{m}_\ell^2}{\hat{s}} \right) \lambda(1, \hat{s}, \hat{m}_\rho^2) \end{aligned} \quad (4.14)$$

plugging in the expressions of differential decay rate of $\bar{B} \rightarrow \rho\ell^+\ell^-$ and $B \rightarrow \bar{\rho}\ell^+\ell^-$ given by eqn(4.10) and eqn(4.13) respectively we can get the expression of partial width CP asymmetry :

$$A_{CP}(\hat{s}) = \frac{-2 \text{Im}\lambda_u \Delta_\rho}{\Sigma_\rho + 2 \text{Im}\lambda_u \Delta_\rho} \quad (4.15)$$

with Σ_ρ and Δ_ρ as given in eqn(4.11) and eqn(4.14).

Another measure of CP violation could be the sum of the FB asymmetries of $\bar{B} \rightarrow \rho\ell^+\ell^-$ and $B \rightarrow \bar{\rho}\ell^+\ell^-$. One can calculate this by use of eqn(4.12) for $\bar{B} \rightarrow \rho\ell^+\ell^-$ and for $B \rightarrow \bar{\rho}\ell^+\ell^-$ by making appropriate change in the expression of C_9^{eff} . The final expression is :

$$\begin{aligned} \delta_{FB} = & A_{FB}(\bar{B} \rightarrow \rho\ell^+\ell^-) + A_{FB}(B \rightarrow \bar{\rho}\ell^+\ell^-) \\ = & -6 \lambda^{1/2}(1, \hat{s}, \hat{m}_\rho^2) \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \text{Im}\lambda_u \frac{\left[\text{Im}\xi_2 \Gamma_1 - 2 \Delta_\rho \left\{ \text{Re}(C_9^{eff}) \Gamma_1 + \frac{2 C_7^{eff} \hat{m}_b}{\hat{s}} \Gamma_2 \right\} \right]}{\Sigma_\rho (\Sigma_\rho + 4 \text{Im}\lambda_u \Delta_\rho)} \end{aligned} \quad (4.16)$$




FIG. 9: CP violating asymmetry (δ_{CP}) in $\bar{B} \rightarrow \rho \tau^+ \tau^-$ and $B \rightarrow \bar{\rho} \tau^+ \tau^-$. The Wolfenstein parameter we have chosen are : $\rho = -0.07$, $\eta = 0.34$. Other parameters are : mSUGRA $m = 200$, $M = 450$, $A = 0$, $\tan \beta = 35$, $\text{sgn}(\mu)$ is taken to be +ve for rSUGRA $m_A = 306$. All masses are in GeV

with Γ_1 and Γ_2 given as :

$$\Gamma_1 = 4\hat{s}A_1(s)V(s)C_{10} - \frac{\hat{m}_\ell(1 - \hat{m}_\rho^2 - \hat{s})(1 + \hat{m}_\rho)}{2\hat{m}_\rho\hat{m}_b}A_0(s)A_1(s)C_{Q_1} + \frac{\hat{m}_\ell\lambda(1, \hat{s}, \hat{m}_\rho^2)}{\hat{m}_\rho\hat{m}_b(1 + \hat{m}_\rho)}A_0(s)A_2(s)C_{Q_1} \quad (4.17)$$

$$\Gamma_2 = 4\hat{s}T_2(s)V(s)(1 - \hat{m}_\rho)C_{10} + 4\hat{s}(1 + \hat{m}_\rho)A_1(s)T_1C_{10} - \frac{(1 - \hat{m}_\rho^2 - \hat{s})(1 - \hat{m}_\rho^2)\hat{m}_\ell}{\hat{m}_\rho\hat{m}_b}A_0(s)T_2(s)C_{Q_1} \\ + \frac{2\lambda(1, \hat{s}, \hat{m}_\rho^2)\hat{m}_\ell}{\hat{m}_\rho\hat{m}_b}A_0(s) \left(T_2(s) + \frac{\hat{s}}{1 - \hat{m}_\rho^2}T_3(s) \right) C_{Q_1} \quad (4.18)$$

B. Polarization asymmetries

Finally we calculate the three polarization asymmetries namely longitudinal, transverse and normal, for $\bar{B} \rightarrow \rho \ell^+ \ell^-$. Longitudinal Polarization asymmetry (P_L) is :

$$P_L = \left\{ 24 \text{Re}(A^*B) (1 - \hat{m}_\rho^2 - \hat{s})\hat{s} \left(-1 + \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \right) + 4 m_B^2 \lambda(1, \hat{s}, \hat{m}_\rho^2)\hat{s} \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \text{Re}(A^*D) \right. \\ + \frac{1}{\hat{m}_\rho^2} \left(3 + \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \right) \left[2 \text{Re}(B^*E)(1 + \hat{m}_\rho^4 + 2\hat{m}_\rho^2\hat{s} + \hat{s}^2 - 2(\hat{m}_\rho^2 + \hat{s})) \right. \\ + m_B^2 \text{Re}(C^*E)(1 - 3(\hat{m}_\rho^2 + \hat{s}) - (\hat{m}_\rho^2 - \hat{s})^2(\hat{m}_\rho^2 + \hat{s}) + (3\hat{m}_\rho^4 + 2\hat{m}_\rho^2\hat{s} + 3\hat{s}^2)) \left. \right] \\ + \frac{1}{\hat{m}_\rho^2} \left(\text{Re}(B^*F)(1 - \hat{m}_\rho^2 - \hat{s}) + \text{Re}(C^*F)m_B^2 \lambda(1, \hat{s}, \hat{m}_\rho^2) \right) \left[\left(3 + \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \right) (1 + \hat{m}_\rho^2(\hat{m}_\rho^2 - \hat{s}) - 2\hat{m}_\rho^2) \right. \\ + \left. \left(3 - 7\sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \right) \hat{s}(\hat{m}_\rho^2 - \hat{s}) - 8\hat{s}\sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} \right] \left. \right\} / \Sigma_\rho \quad (4.19)$$




FIG. 10: Longitudinal Polarization asymmetry (P_L) in $\bar{B} \rightarrow \rho\tau^+\tau^-$. The Wolfenstein parameter we have chosen are : $\rho = -0.07$, $\eta = 0.34$. Other parameters are : mSUGRA $m = 200$, $M = 450$, $A = 0$, $\tan\beta = 35$, $\text{sgn}(\mu)$ is taken to be +ve for rSUGRA $m_A = 306$. All masses are in GeV

Normal Polarization asymmetry (P_N) is :

$$P_N = \lambda^{1/2}(1, \hat{s}, \hat{m}_\rho^2) \sqrt{(\hat{s} - 4\hat{m}_\ell^2)} \pi \left[2\text{Im}(E^*F) \frac{1 + \hat{m}_\rho^2 - \hat{s}}{\hat{m}_\rho^2} + 2\text{Im}(A^*E + B^*D) \right. \\ \left. + \frac{1}{4\hat{m}_\rho^2} \{ 2(1 - \hat{m}_\rho^2 - \hat{s})\text{Im}(G^*B) + m_B^2 \lambda(1, \hat{s}, \hat{m}_\rho^2) \text{Im}(G^*C) \} \right] \quad (4.20)$$

finally Transverse Polarization asymmetry (P_T) is :

$$P_T = \lambda^{1/2}(1, \hat{s}, \hat{m}_\rho^2) \sqrt{\hat{s}} \hat{m}_\ell \pi \left[-4 \text{Re}(A^*B) \right. \\ \left. + \frac{1}{4\hat{m}_\rho^2 \hat{s}} \{ 2(2(1 - \hat{m}_\rho^2 - \hat{s})\text{Re}(B^*E) + m_B^2 \lambda(1, \hat{s}, \hat{m}_\rho^2) \text{Re}(C^*E)) \} \right] \quad (4.21)$$

V. RESULTS AND DISCUSSION

We have performed the numerical analysis of all the kinematical variables which we have evaluated in Section III and Section IV.

For our numerical analysis we would be using the Minimal Supersymmetric Standard Model (MSSM), which is the simplest (and the one having the least number of parameters) extension of the SM. Actually MSSM itself has fairly large number of parameters which makes it difficult to do phenomenology with it. We therefore resort to models which reduces the large parameter space of MSSM to a manageable level. The models which are around includes minimal Supergravity (mSUGRA), no-scale, dilaton etc. models. For our analysis we will be going to use Supergravity

(SUGRA) model. The basic features of all these models is that they assume some sort of unification of the parameters at some unifying scale. In the numerical analysis of SUGRA models (both mSUGRA and rSUGRA) the parameters we have choosen satisfies the radiative electroweak symmetry breaking condition.

In mSUGRA model unification of all the scalar masses, all gaugino masses and all coupling constants unification is assumed at GUT scale. So effectively we would be left with five parameters (over the SM parameters) at GUT scale they are : m (unified mass of all the scalars), M (unified mass of all the gauginos), A (unified trilinear coupling constants), $\tan\beta$ (the ratio of vacuum expectation values of the two Higgs doublets) and finally the $sgn(\mu)$ ⁵. As emphasized in many works [4, 6, 10] the universality of the scalar masses is not a necessary requirement and one can relax this. The only constraint for this relaxation is $K^0 - \bar{K}^0$ mixing. To suppress this mixing it is sufficient to give unified scalar mass to all the squarks but the Higgs sector can be given a different unified mass. We will be going to explore sort of SUGRA model also, which we will be going to call as rSUGRA (or simply SUGRA) model. In this sort of model we will take mass of the pseudo-scalar Higgs to be another parameter. For our MSSM parameter space analysis we will be going to take 95% CL bound [21] :

$$2 \times 10^{-4} < Br(B \rightarrow X_s \gamma) < 4.5 \times 10^{-4}$$

which is in agreement with CLEO and ALPEH results. As we are more interested in finding out the effects of NHBs and as well emphasized in literature that these effects becomes more profound when the final state leptons are τ [18]. So we will be going to take the final state leptons to be τ here.




FIG. 11: Normal Polarization asymmetry (P_N) in $\bar{B} \rightarrow \rho\tau^+\tau^-$. The Wolfenstein parameter we have choosen are : $\rho = -0.07$, $\eta = 0.34$. Other parameters are : mSUGRA $m = 200$, $M = 450$, $A = 0$, $\tan\beta = 35$, $sgn(\mu)$ is taken to be +ve for rSUGRA $m_A = 306$. All masses are in GeV

⁵ our sign of convention of μ is such that μ enters the chargino mass matrix with positive sign




FIG. 12: Transverse Polarization asymmetry (P_T) in $\bar{B} \rightarrow \rho\tau^+\tau^-$. The Wolfenstein parameter we have chosen are : $\rho = -0.07$, $\eta = 0.34$. Other parameters are : mSUGRA $m = 200$, $M = 450$, $A = 0$, $\tan\beta = 35$, $\text{sgn}(\mu)$ is taken to be +ve for rSUGRA $m_A = 306$. All masses are in GeV

Our results are summarized in Figures 1 - 12, where the spikes in the distributions are because of the charm resonances as given in eqn.(2.2). For $B \rightarrow \pi\tau^+\tau^-$, in Figure(1) we have plotted the variation of Branching ratio with the scaled cm energy of the dileptons. As we can see from the graphs the deviation from the respective SM values is fairly large for almost the whole region of invariant dilepton mass. The deviation is more profound for SUGRA model than mSUGRA model. In Figure(2) we have plotted the FB asymmetry for the transition. As has already been noted in earlier works [8] that in SM the FB asymmetry vanishes. But if we consider SUSY then one can have a finite value of FB asymmetry. So any observation of FB asymmetry in this decay mode ($B \rightarrow \pi\tau^+\tau^-$) should be a clear signal of new physics. In Figure(3) we have given the estimates of the CP violating partial width asymmetry. As expected from the result of eqn(3.15) the new Wilson coefficients doesn't give any contributions to the numerator of the asymmetry but the denominator (which essentially is the decay rate) gets contributions from NHBs and hence the results that NHB effects actually lowers the SM estimates of the CP violating partial width asymmetry. The reduction is more for SUGRA model where all the scalar masses are not unified, in this case one can take Higgs mass to be also a parameter ⁶. The new Wilson coefficients C_{Q_1} and C_{Q_2} crucially depends on Higgs mass and if Higgs mass is low their value is high. In SUGRA model one can have relatively lower Higgs mass and hence high value of new Wilsons and hence high partial decay rate which effectively reduces the CP violating partial width asymmetry. But there exists another measure of CP violation which is the sum of the FB asymmetries of $\bar{B} \rightarrow \pi\ell^+\ell^-$ and $B \rightarrow \bar{\pi}\ell^+\ell^-$. This is a sort of measurement which can be done in an environment having equal numbers of B and $\bar{B}$ pairs and as argued earlier [13] doesn't require any tagging. The important point over here is that FB asymmetry in this decay is zero in SM and hence the sum of the FB asymmetries of $\bar{B} \rightarrow \pi\tau^+\tau^-$ and $B \rightarrow \bar{\pi}\tau^+\tau^-$. So the parameter δ_{FB} (which we have introduced in eqn(3.17) is zero. But if we consider SUSY then this parameter can have a finite value.

⁶ here we have taken pseudo-scalar Higgs mass to be a parameter and all the rest Higgs masses can be evaluated in terms of this

TABLE I: Integrated kinematical variables for $B \rightarrow \pi\tau^+\tau^-$. The parameters of mSUGRA and rSUGRA are same as given in Figures(1) - (5)

Variable	SM	mSUGRA	rSUGRA
$d\Gamma/d\hat{s} \times 10^8$	2.6	3.43	5.04
$A_{FB} \times 10$	0	- 0.224	- 0.228
$A_{CP} \times 10^2$	0.51	0.2	0.1
$\delta_{FB} \times 10^3$	0	- 0.6	- 0.9
$P_N \times 10^2$	0	0.96	0.99

In fact as we have shown in Figure(4) δ_{FB} is more for SUGRA model then mSUGRA model (which is contrary to the partial width CP asymmetry). So this quantity could also turn out to be important probe for new physics. In Figure(5) we have plotted the normal polarization asymmetry of the final state lepton ℓ^- . As we can see from the expression of eqn(3.21) the value of P_N is zero in SM. So the observation of non-zero P_N could also be interpreted as signal of some new physics ⁷. Rest two polarization asymmetries the longitudinal (P_L) and transverse (P_T) vanishes with or without NHBs.

In Figure(6) we have plotted the Branching ratio of $\bar{B} \rightarrow \rho\tau^+\tau^-$ with scaled invariant mass of dileptons. As we can see there is a fairly large deviation from the SM value. In Figure(7) we have plotted the variation of FB asymmetry with $\hat{s}$ again one can observe the variation of mSUGRA and SUGRA results from SM values. Both the partial decay rate and FB asymmetry increase as compared to SM values in both mSUGRA and SUGRA models. In Figure(8) we have plotted the CP violating partial width asymmetry. As we can see that here the predictions of mSUGRA and SUGRA model gets decreased as compared to the SM value. The reason is the same as explained for $B \rightarrow \pi\tau^+\tau^-$ transition. But again here if we look at the CP violation through FB asymmetry (Figure(9)) and we have the same effect as there in $B \rightarrow \pi\tau^+\tau^-$ the SUGRA models increases values of as compared to SM values. Although here the SM values of FB asymmetry as well as δ_{FB} is not zero but still SUGRA models gives a enhancement of more than a order (, i.e. ,about a factor of 10) for almost whole region of the invariant dileptonic mass. In Figures(10),(11),(12) we have plotted the Longitudinal (P_L), Normal (P_N) and Transverse (P_T) polarization respectively respectively. All the three shows variation from SM values but the general tend is that all these polarization asymmetries tends to decrease in SUGRA models as compared to SM values for almost whole region of invariant mass. For all the plots we have taken the variation of all the kinematical variables with the dileptonic invariant mass because the variation with respect to (wrt) invariant mass has more information than the result which we get after integrating over the invariant mass and also the variation of kinematical variables wrt invariant mass are in principle accessible experimentally. In future B-factories (like Tevatron and LHCb) more than 10^{11} , $b\bar{b}$ pairs are expected to be produced [21]; this is many orders more than the projected yeild at the e^+e^- B factories, so these processes can be observed and some of the measurable quantities in these processes can be estimated. These processes ($B \rightarrow \pi\ell^+\ell^-$ and $B \rightarrow \rho\ell^+\ell^-$) are useful because they are relatively clean (both theoretically and experimentally). Also if there is no new source of CP violation (except CKM phase) then dileptonic decays $B \rightarrow \pi\ell^+\ell^-$ and $B \rightarrow \rho\ell^+\ell^-$ should be the first one where CP violation would be observed. As we can see from the Tables I, II that the kinematical variables of $B \rightarrow \rho\tau^+\tau^-$ looks to be more promising because of their magnitude. But in $B \rightarrow \pi\tau^+\tau^-$ there are some distributions like FB-asymmetry,

⁷ P_N is a T-odd observable which comes because of the non-hermiticity of the effective Hamiltonian, associated with the real $c\bar{c}$ intermediate states, so it can't be taken as a measure of CP violation

TABLE II: Integrated kinematical variables for $B \rightarrow \rho\tau^+\tau^-$. The parameters of mSUGRA and rSUGRA are same as given in Figures(6) - (12)

Variable	SM	mSUGRA	rSUGRA
$d\Gamma/d\hat{s} \times 10^8$	3.9	4.4	5.0
$A_{FB} \times 10$	-0.72	- 0.60	- 0.32
$A_{CP} \times 10$	0.13	0.09	0.04
δ_{FB}	0.0003	- 0.11	- 0.18
P_L	0.109	0.0924	0.055
$P_N \times 10$	0.16	0.14	0.1
P_T	0.17	0.14	0.07

δ_{FB} and P_N which vanishes in SM but gives finite (although small) values with SUSY. This point has already been noted about FB-asymmetry in many other earlier works [8]. The same phenomena occurs for δ_{FB} and P_N .

Finally our conclusions regarding SUSY effects over a wide range of SUSY parameters can be summarized as :

1. **Branching Ratios** : The branching ratios for both $B \rightarrow \pi\tau^+\tau^-$ and $B \rightarrow \rho\tau^+\tau^-$ shows large deviation from respective SM values for almost whole region of the invariant mass.
2. **FB asymmetry** : FB asymmetry for $B \rightarrow \pi\ell^+\ell^-$ vanishes within SM. A non-vanishing FB asymmetry for $B \rightarrow \pi\ell^+\ell^-$ clearly gives indications of some new physics. FB asymmetry for $B \rightarrow \rho\ell^+\ell^-$ shows significant increase in mSUGRA and rSUGRA models as compared to SM values.
3. **CP violating partial width asymmetry (A_{CP})** : The predictions of mSUGRA and SUGRA models is to reduce this asymmetry for both $B \rightarrow \pi\tau^+\tau^-$ and $B \rightarrow \rho\tau^+\tau^-$ as compared to SM values.
4. **CP violation from FB asymmetry (δ_{FB})** : This observable vanishes in SM for $B \rightarrow \pi\tau^+\tau^-$ a non-vanishing value of this clearly indicates new physics effects. For $B \rightarrow \rho\tau^+\tau^-$ the SM value prediction is very low, both SUGRA and mSUGRA can give enhancement of over a order almost for whole region of scaled dilepton invariant mass.
5. **Polarization asymmetries** : For $B \rightarrow \pi\tau^+\tau^-$ all the three polarization asymmetries (longitudinal, normal and transverse) vanishes in SM. If we include NHB effects then although longitudinal and transverse still remains zero the normal polarization asymmetry becomes non-zero. So observation of normal polarization asymmetry can still be regarded as evidence for new physics. For $B \rightarrow \rho\ell^+\ell^-$ all the three polarization asymmetries decreases, with respect to their respective SM values, by switching on the NHB effects.

The observation of the decay modes $B \rightarrow \pi\ell^+\ell^-$ and $B \rightarrow \rho\ell^+\ell^-$ can be expected to be a very useful tool for the search of new physics effects as well as for the measurement of the CP violating parameters of CKM matrix.

Acknowledgments

This work was supported under SERC scheme of Department of Science and Technology (DST), India.

APPENDIX A: INPUT PARAMETERS

$$\begin{aligned}
m_u &= m_d = 10 \text{ MeV} \\
m_b &= 4.8 \text{ GeV} , \quad m_c = 1.4 \text{ GeV} \quad m_t = 176 \text{ GeV} \\
m_B &= 5.26 \text{ GeV} , \quad m_\pi = 0.135 \text{ GeV} , \quad m_\rho = 0.768 \text{ GeV} \\
|V_{tb}V_{td}^*| &= 0.011 , \quad \alpha = \frac{1}{129} , \quad G_F = 1.17 \times 10^{-5} \text{ GeV}^{-2} \\
m_\tau &= 1.77 \text{ GeV} , \quad \tau_B = 1.54 \times 10^{-12} \text{ s} \\
\text{Wolfenstein parameters : } \rho &= -0.07 , \quad \eta = 0.34
\end{aligned}$$

APPENDIX B: FORM FACTORS FOR $B \rightarrow \pi$ TRANSITION

We will be going to use the form factors given by Coleangelo et. al. [23] :

$$\langle \pi(p_\pi) | \bar{d} \gamma_\mu P_{L,R} b | B(p_B) \rangle = \frac{1}{2} \left\{ (2p_B - q)_\mu F_1(q^2) + \frac{m_B^2 - m_\pi^2}{q^2} q_\mu (F_0(q^2) - F_1(q^2)) \right\} \quad (\text{B1})$$

$$\langle \pi(p_\pi) | \bar{d} i \sigma_{\mu\nu} q^\nu P_{L,R} b | B(p_B) \rangle = \frac{1}{2} \left\{ (2p_B - q)_\mu - (m_B^2 - m_\pi^2) q_\mu \right\} \frac{F_T(q^2)}{m_B + m_\pi} \quad (\text{B2})$$

To get the matrix element for scalar current we multiply eqn.(B1) by q_μ we get

$$\langle \pi(p_\pi) | \bar{d} P_R b | B(p_B) \rangle = \frac{1}{2m_b} (m_B^2 - m_\pi^2) F_0(q^2) \quad (\text{B3})$$

where we have neglected the mass of d-quark.

The definition of the Form factors F_0 , F_1 and F_T are ⁸ (q^2 in the units of GeV^2) :

$$\begin{aligned}
F_0(q^2) &= \frac{F_0(0)}{1 - q^2/7^2} \\
F_1(q^2) &= \frac{F_1(0)}{1 - q^2/5.3^2} \\
F_T(q^2) &= \frac{F_T(0)}{(1 - q^2/7^2)(1 - q^2/5.3^2)} \\
\tilde{F}_T(q^2) &= \frac{F_T(q^2)}{(m_B + m_\pi)} m_b
\end{aligned} \quad (\text{B4})$$

with $F_0(0) = 0$, $F_1(0) = 0.25$ and $F_T(0) = -0.14$

APPENDIX C: FORM FACTORS FOR $B \rightarrow \rho$ TRANSITION

For $B \rightarrow \rho$ transition we will be going to use the form factors given by Colangelo et. al. [23] :

$$\begin{aligned}
\langle \rho(p_\rho) | \bar{d} \gamma_\mu P_L b | \bar{B}(p_B) \rangle &= i \epsilon_{\mu\nu\alpha\beta} \epsilon^{\nu*} p_B^\alpha q^\beta \frac{V(q^2)}{m_B + m_\rho} - \frac{1}{2} \left\{ \epsilon_\mu (m_B + m_\rho) A_1(q^2) \right. \\
&\quad \left. - (\epsilon^* \cdot q) (2p_B - q)_\mu \frac{A_2(q^2)}{m_B + m_\rho} - \frac{2m_\rho}{q^2} (\epsilon^* \cdot q) [A_3(q^2) - A_0(q^2)] \right\}
\end{aligned} \quad (\text{C1})$$

⁸ All the three F_0 , F_1 and F_T are not independent. F_T can be related to F_0 and F_1 by equation of motion and the relationship works out to be $F_T = (m_B + m_\pi) m_b \frac{F_0 - F_1}{q^2}$

$$\begin{aligned} \langle \rho(p_\rho) | \bar{d} i \sigma_{\mu\nu} q^\nu P_{R,L} b | \bar{B}(p_B) \rangle &= -2i \epsilon_{\mu\nu\alpha\beta} \epsilon^{\nu*} p_B^\alpha q^\beta T_1(q^2) \pm [\epsilon_\mu^* (m_B^2 - m_\rho^2) - (\epsilon^* \cdot q) (2 p_B - q)_\mu] T_2(q^2) \\ &\pm (\epsilon^* \cdot q) \left[q_\mu - \frac{q^2}{m_B^2 - m_\rho^2} (2 p_B - q)_\mu \right] T_3(q^2) \end{aligned} \quad (C2)$$

where A_3 can be written in terms of A_1 and A_2 , i.e. ,

$$A_3(q^2) = \frac{m_B + m_\rho}{2m_\rho} A_1(q^2) - \frac{m_B - m_\rho}{2m_\rho} A_2(q^2) \quad (C3)$$

In above equations ϵ is the polarization vector of ρ and $q = p_B - p_\rho$ is the momentum transfer.

To get the matrix element for scalar (or pseudoscalar) current we multiply both sides of eqn.(C1) by q^μ . On simplifying we get :

$$\langle \rho(p_\rho) | \bar{d} P_R b | B(p_B) \rangle = - \frac{m_\rho}{m_b} (\epsilon^* \cdot q) A_0(q^2) \quad (C4)$$

The definition of the form factors is ⁹ (q^2 in units of GeV^2) :

$$\begin{aligned} V(q^2) &= \frac{V(0)}{1 - q^2/5^2} \\ A_1(q^2) &= A_1(0) (1 - 0.023 q^2) \\ A_2(q^2) &= A_2(0) (1 + 0.034 q^2) \\ A_0(q^2) &= \frac{A_3(0)}{1 - q^2/4.8^2} \\ T_1(q^2) &= \frac{T_1(0)}{1 - q^2/5.3^2} \\ T_2(q^2) &= T_2(0) (1 - 0.02 q^2) \\ T_3(q^2) &= T_3(0) (1 + 0.005 q^2) \end{aligned} \quad (C5)$$

with $V(0) = 0.47$, $A_1(0) = 0.37$, $A_2(0) = 0.4$, $A_0(0) = 0.3$, $T_1(0) = 0.19$, $T_2(0) = 0.19$, $T_3(0) = -0.7$.

-
- [1] A. Ali, arXiv:hep-ph/9709507,
G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996) [arXiv:hep-ph/9512380].
- [2] E. O. Iltan and G. Turan, Phys. Rev. **D 61**, 034010 (2000)[arXiv:hep-ph/9906502] ,
G. Erkol and G. Turan, hep-ph/0112115
- [3] W. Skiba and J. Kalinowski, Nucl. Phys. **B 404**, 3 (1993),
H. E. Logan and U. Nierste, Nucl. Phys. **B 586**, 39 (2000)[arXiv:hep-ph/0004139].
Y. Dai, C. Huang and H. Huang, Phys. Lett. **B 390**, 257 (1997)[arXiv:hep-ph/9607389],
- [4] S. R. Choudhury and N. Gaur, Phys. Lett. **B 451**, 86 (1999)[arXiv:hep-ph/9810307].
- [5] Z. Xiong and J. M. Yang, Nucl. Phys. **B 628**, 193 (2002)[arXiv:hep-ph/0105260].
C. Bobeth, A. J. Buras, F. Kruger and J. Urban, Nucl. Phys. **B 630**, 87 (2002) [arXiv:hep-ph/0112305].
C. Huang, W. Liao and Q. Yan, Phys. Rev. **D 59**, 011701 (1999)[arXiv:hep-ph/9803460],
- [6] S. Rai Choudhury, A. Gupta and N. Gaur, Phys. Rev. **D 60**, 115004 (1999)[arXiv:hep-ph/9902355].
S. R. Choudhury, N. Gaur and A. Gupta, Phys. Lett. **B 482**, 383 (2000)[hep-ph/9909258].

⁹ again here also T_3 can be related to A_3 and A_0 by EOM and relationship is : $T_3 = m_B m_b \frac{A_3 - A_0}{q^2}$

- [7] S. Rai Choudhury, Naveen Gaur and Namit Mahajan, Phys. Rev. **D 66**, 054003 (2002)[arXiv:hep-ph/0203041]
S. Rai Choudhury and Naveen Gaur arXiv:hep-ph/0205076
- [8] D. A. Demir, K. A. Olive and M. B. Voloshin Phys. Rev. **D 66**, 034015 (2002)[arXiv : hep-ph/0204119] ;
C. Bobeth, T. Ewerth, F. Kruger and J. Urban , Phys. Rev. **D 64**, 074014 (2001) , [arXiv : hep-ph/0104284].
G. Erkol and G. Turan arXiv : hep-ph/0201055
- [9] E. O. Iltan , Int. J. Mod. Phys. **A 14**, 4365 (1999), [arXiv : hep-ph/9807256] ;
T. M. Aliev and M. Savci , Phys. Rev. **D 60**, 014005 (1999), [arXiv : hep-ph/9812272].
- [10] T. Goto, Y. Okada, Y. Shimizu and M. Tanaka, Phys. Rev. **D 55**, 4273 (1997)[arXiv:hep-ph/9609512],
T. Goto, Y. Okada and Y. Shimizu, Phys. Rev. **D 58**, 094006 (1998)[arXiv:hep-ph/9804294].
J. R. Ellis, K. A. Olive and Y. Santoso arXiv : hep-ph/0204192
- [11] P. L. Cho, M. Misiak and D. Wyler, Phys. Rev. **D 54**, 3329 (1996)[arXiv:hep-ph/9601360].
J. L. Hewett and J. D. Wells, Phys. Rev. **D 55**, 5549 (1997)[arXiv:hep-ph/9610323].
- [12] D. S. Du and M. Z. Yang, Phys. Rev. **D 54**, 882 (1996) [arXiv:hep-ph/9510267].
T. M. Aliev, D. A. Demir, E. Iltan and N. K. Pak, Phys. Rev. **D 54**, 851 (1996)[arXiv:hep-ph/9511352].
- [13] S. Rai Choudhury, Phys. Rev. **D 56**, 6028 (1997), [arXiv : hep-ph/9706313].
- [14] F. Kruger and L. M. Sehgal, Phys. Rev. **D 56**, 5452 (1997)[Erratum-ibid. **D 60**, 099905 (1997)] [arXiv:hep-ph/9706247].
- [15] S. Fukae, C. S. Kim and T. Yoshikawa, Phys. Rev. **D 61**, 074015 (2000)[arXiv:hep-ph/9908229].
T. M. Aliev, M. K. Cakmak and M. Savci, Nucl. Phys. B **607**, 305 (2001) [arXiv:hep-ph/0009133].
T. M. Aliev, M. K. Cakmak, A. Ozpineci and M. Savci, Phys. Rev. **D 64**, 055007 (2001) [arXiv:hep-ph/0103039].
- [16] F. Krüger and L. M. Sehgal, Phys. Lett. **B 380**, 199 (1996)[arXiv:hep-ph/9603237],
J. L. Hewett, Phys. Rev. **D 53**, 4964 (1996)[arXiv:hep-ph/9506289].
- [17] F. Kruger and L. M. Sehgal, Phys. Rev. **D 55**, 2799 (1997)[arXiv:hep-ph/9608361].
- [18] Y. Grossman, Z. Ligeti and E. Nardi, Phys. Rev. **D 55**, 2768 (1997)[arXiv:hep-ph/9607473],
- [19] B. Grinstein, M. J. Savage and M. B. Wise, Nucl. Phys. **B 319**, 271 (1989),
A. J. Buras and M. Münz, Phys. Rev. **D 52**, 186 (1995)[arXiv:hep-ph/9501281].
- [20] A. Ali, T. Mannel and T. Morozumi, Phys. Lett. **B 273**, 505 (1991).
C. S. Lim, T. Morozumi and A. I. Sanda, Phys. Lett. **B 218**, 343 (1989).
N. G. Deshpande, J. Trampetic and K. Panose, Phys. Rev. **D 39**, 1461 (1989).
P. J. O'Donnell and H. K. Tung, Phys. Rev. **D 43**, 2067 (1991) .
- [21] B Physics at Tevatron : Run II & Beyond , K. Anikeev et. al. arXiv : hep-ph/0201071,
CLEO collaboration, T.E.Coan et. al. Phys. Rev. Lett. **84**, 5283 (2000)[arXiv:hep-ex/9908022] ,
ALEPH Collaboration, R.Barate et. al. Phys. Lett. **B 429**, 169 (1998).
- [22] L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1983.)
- [23] P. Coleangelo, F. De Fazio, P. Santorelli and E. Scrimieri Phys. Rev. **D 53**, 3672 (1996)